



# ***In-vitro* Antioxidant Capacity, Phytochemical Characterisation, Toxic and Functional Properties of African Yam Bean (*Sphenostylis stenocarpa*) Seed-Enriched Cassava (*Manihot esculenta*) Product (*Pupuru*)**

Folayemi Janet Isaac-Bamgboye<sup>1\*</sup>, Victor Ndigwe Enujiugha<sup>1</sup>  
and Matthew Olusola Oluwamukomi<sup>1</sup>

<sup>1</sup>Department of Food Science and Technology, Federal University of Technology,  
P.M.B. 747, Akure, Nigeria.

## **Authors' contributions**

*This work was carried out in collaboration among all authors. Author FJIB designed the study, carried out the chemical analysis, performed the statistical analysis, managed the literature searches, the protocol and wrote the first draft of the manuscript. Authors VNE and MOO designed and supervised the study. All authors read and approved the final manuscript.*

## **Article Information**

DOI: 10.9734/EJNFS/2020/v12i330212

### Editor(s):

(1) Dr. Rasha Mousa Ahmed Mousa, University of Jeddah, Saudi Arabia.

### Reviewers:

(1) Bagiu Radu-Vasile, Victor Babeş University of Medicine and Pharmacy, Timișoara, Romania.

(2) Tugay Ayasan, Osmaniye Korkut Ata University, Turkey.

Complete Peer review History: <http://www.sdiarticle4.com/review-history/55044>

**Original Research Article**

**Received 07 February 2020**

**Accepted 12 April 2020**

**Published 05 May 2020**

## **ABSTRACT**

**Aims:** This study aimed at determining the *In-vitro* antioxidant capacity, characterise phytochemical constituents, assess toxic and functional properties of African yam bean (*Sphenostylis stenocarpa*) seed-enriched Cassava product (*Pupuru*) flour blends using standard methods.

**Methodology:** *Pupuru* flour blends were produced from spontaneously-fermented cassava tubers substituted with African yam bean (*Sphenostylis stenocarpa*) seed (AYBS) (5% (EP5), 10% (EP10) and 15% (EP15), before toasting, cooling, milling, sieving and packaging. A commercial sample (CP) with 100% cassava and another produced in this study, were used as controls.

\*Corresponding author: Email: [yemistyle@gmail.com](mailto:yemistyle@gmail.com);

**Results:** *In-vitro* 2,2-Azino-bis (3-ethylbenzothiazoline-6-sulphonic acid) (ABTS) and 1,1-Diphenyl-2-picrylhydrazyl (DPPH) free radical scavenging abilities, total flavonoid content (TFC), ferric reducing antioxidant power (FRAP) and total phenol content (TPC) increased significantly ( $P = .05$ ) as AYBS enrichment levels increased. The commercial and laboratory control samples showed no significant difference ( $P = .05$ ) in all the antioxidants analysed except DPPH: ABTS (7.61- 12.27%); DPPH (26.34-48.26%); TFC (0.10-0.25 mg CAE/g); FRAP (0.81-2.36 (mg/g) and TPC (15.74-24.15 mg GAE/g). All the phytochemicals except tannins increased significantly ( $P = .05$ ) as levels of enrichment with AYBS increased. Tannins, phytates, saponins, alkaloids and oxalates were 1.46 -2.87 (mg/g); 0.85-1.40 mg/100 g; 4.18-13.27 mg/g; 24.89-29.05 mg/g and 1.71-3.23 mg/g, respectively. The toxic constituent revealed that all the samples contained significantly different ( $P = .05$ ) cyanide ranging from 0.87-2.51 mg/kg which reduced as AYBS level of inclusion increased. The functional properties of the samples were significantly ( $P = .05$ ) enhanced with AYBS enrichment.

**Conclusion:** Utilisation of AYBS to enrich *Pupuru* increased its *In-vitro* antioxidant capacity and phytochemical constituents, reduced the toxic cyanide content enhanced the functional properties, hence, its suitability as a nutraceutical to delay the ageing process and prevent cardiovascular diseases.

**Keywords:** African yam bean (*Sphenostylis stenocarpa*) seeds; cassava; antioxidant; cyanide, phytochemicals; functional properties; nutraceuticals.

## 1. INTRODUCTION

A predictable upsurge in the global request for food in the next few decades is looming owing to continuous population and consumption growth [1]. The last five decades have experienced remarkable growth in food production, resulting in a marked decline in the percentage of the hungry people, globally, despite a double increase in the total population [2,3]. The natural antioxidant content of food products has become the focus of most studies in recent years owing to its involvement in defence against diseases [4]. Also, there has been a recent growing indication to propose that countless age-associated human diseases such as cancer, arthritis, immune system degeneration, brain dysfunction, heart diseases and cataracts are the consequences of cellular impairment by free radicals and that antioxidants in foods could play an imperative role in the prevention of such diseases [5,6,7]. The public attention in natural antioxidants has been much driven and steered towards an all-encompassing search for effective, yet natural antioxidants in foods [6-10]. The capacity of some inherent bioactive compounds in food products, such as African yam bean (*Sphenostylis stenocarpa*) seed (AYBS), to work as antioxidants has been documented, resulting to the impending benefits of consuming foods that are rich in such compounds [7,11-13]. The phenolic extracts from AYBS was reported to have exhibited antioxidant power and the ability to scavenge free radicals [12] and the researchers established that

determination of antioxidant activity of the AYB extracts was a function of the use of appropriate solvent and concentration of extracts (the higher the concentration of extracts, the higher the capacity to scavenge free radicals).

*Pupuru* is a stiff dough meal traditionally prepared from fermented, smoke-dried cassava (*Manihot esculenta* Crantz) roots and consumed by the riverine dwellers of the South-Western States in Nigeria. Cassava root is primarily starchy, energy-dense, with about 80 to 90% carbohydrate on a dry weight basis, with toxic and anti-nutritional components like cyanide and phytate [14,15]. There seemed to be scarce studies on the antioxidant activity, toxic, anti-nutritional and functional properties of AYBS-enriched *Pupuru* flour blends. Because of the anticipated shortage of foods, globally, this study aimed at determining the effect of AYBS enrichment of cassava on the *in-vitro* antioxidant activities, phytochemical constituents, toxic and functional properties of the developed *Pupuru* flour blends.

## 2. MATERIALS AND METHODS

### 2.1 Materials

Fresh cassava (*Manihot esculenta*, Crantz) (TME 0581) tubers were sourced from the research farm of the Federal University of Technology, Akure, Ondo State. African Yam Bean (AYB) (*Sphenostylis stenocarpa*) seeds (TSs 091) were

from a farm in Efon Alaaye, Ekiti State and characterised by the International Institute of Tropical Agriculture (IITA), Ibadan, Oyo State. Reagents and chemicals of analytical grade were purchased from a local chemical store in Akure, Ondo State.

## 2.2 Methods

### 2.2.1 Pre-processing of African Yam Bean (AYB) (*Sphenostylis stenocarpa*) seeds

African yam bean seeds (AYBS) were pre-processed into flour using the method described by Oluwamukomi and Akinlabi [16], with slight modifications. The seeds were manually cleaned to remove dirt, stones, defective seeds, dead insects, and other unwanted materials. The cleaned seeds were washed with and soaked in clean tap water for 24 h, manually dehulled, rinsed, cooked for about 45 min, drained of water, rinsed, dried in an oven at 60°C for 5 h, cooled to about 30±2°C, milled, sieved and the resultant flour was packaged in an airtight container for further uses [16].

### 2.2.2 Production of *Pupuru* flour blends

One kilogram (1 kg) of peeled, chopped and washed cassava tubers were soaked in one litre of water in a sterile plastic container and left to ferment for 96 h. Fibrous materials were manually removed from the dewatered, fermented cassava which was then pressed with a hydraulic jack, for further removal of water before being pulverized. Various percentages (5, 10 and 15) of the African yam bean (*Sphenostylis stenocarpa*) seed (AYBS) flour were homogeneously mixed with the fermented cassava paste before toasting in an open and wide pan until it was dry [16]. The resultant cassava and AYBS blends (Table 1) were cooled to about 30 ± 2°C, milled, sieved and packaged in an airtight container for further analyses. A commercial sample of *Pupuru* flour (CP) with 100% cassava and another prepared during this study (P100) were both used as commercial and laboratory controls, respectively.

**Table 1. Formulation of *Pupuru* flour blends**

Sample code	Formulation ratio AYBS flour: Cassava (%)
P100	0 : 100
EP5	5 : 95
EP10	10 : 90
EP15	15 : 85

AYBS = African yam bean (*Sphenostylis stenocarpa*) seed

## 2.3 In-vitro Antioxidant Capacity Determination

### 2.3.1 Determination of 2, 2 - azino-bis (3-ethylbenthiazoline-6-sulphonic acid) (ABTS) scavenging capacity

The 2, 2 - Azino-bis (3-ethylbenthiazoline-6-sulphonic acid) (ABTS) scavenging ability of each sample was determined according to the method described by Re et al. [17]. The ABTS was generated by reacting an (7 mM) ABTS aqueous solution with K<sub>2</sub>S<sub>2</sub>O<sub>8</sub> (2.45 mM/l, final concentration) in the dark for 16 h and adjusting the absorbance at 734 nm to 0.700 with ethanol. Appropriate dilution of the sample (0.2 ml) was then added to 2.0 ml of the ABTS solution and the absorbance read at 732 nm after 15 m. The Trolox equivalent antioxidant capacity was subsequently calculated as follows:

$$\text{Radical scavenging(\%)} = 100 - [100 \times (A_{\text{sample}} - A_{\text{blank}}) / A_{\text{control}}] \quad (1)$$

Where:

$A_{\text{sample}}$  = the absorbance of the ABTS mixed with the sample,

$A_{\text{control}}$  = the absorbance of the ABTS mixed with water, and

$A_{\text{blank}}$  = the absorbance of sample mixed with water.

### 2.3.2 Determination of 1,1-diphenyl-2-picrylhydrazyl (DPPH) free radical scavenging ability

The free radical scavenging ability of each of the *Pupuru* flour blend extracts against 1,1-diphenyl-2-picrylhydrazyl (DPPH) free radical was evaluated as described by Gyamfi et al. [18]. Briefly, appropriate dilution of the extracts (1 mL) was mixed with 1 mL, 0.4 mM methanolic solution containing DPPH radicals, the mixture was left in the dark for 30 min and the absorbance was taken at 516 nm. The DPPH free radical scavenging ability was subsequently calculated.

$$\text{DPPH (\%)} = \frac{\text{Abs}_{\text{ref}} - \text{Abs}_{\text{standard}}}{\text{Abs}_{\text{ref}}} \times 100 \quad (2)$$

Where,

$\text{Abs}_{\text{ref}}$  = Absorbance of the reference (reacting mixture without the test sample) and,  $\text{Abs}_{\text{sample}}$  =

Absorbance of reacting mixture with the test sample.

### 2.3.3 Determination of Total Flavonoid Content (TFC)

Total Flavonoid Content was determined by aluminium chloride colorimetric assay [19] with slight modification. About 500 µl of methanol was added to 10 ml flask containing 500 µl of aqueous extract. To this 50 µl 10% AlCl<sub>3</sub> and 50 µl of 1 M CH<sub>3</sub>CO<sub>2</sub>K was added, respectively. The total volume was made up to 2500 µl with distilled water. The solution was then incubated at room temperature for 30 min. Absorbance was read against blank at 540 nm with a spectrometer. (JENWAY 6305, United Kingdom). The flavonoid was calculated using quercetin as standard.

$$\text{Total flavonoid content (mg QE/g)} = \frac{\text{Abs.sample} \times \text{Conc.standard (mg/ml)}}{\text{Abs.standard} \times \text{Conc.sample (mg/g)}} \quad (3)$$

Abs<sub>standard</sub> is the absorbance of the solution containing 500 µl quercetin, About 50 µl 10% AlCl<sub>3</sub> and 1 M CH<sub>3</sub>CO<sub>2</sub>K. Blank is the mixture of 500 µl of distilled water, 500 µl of methanol, 50 µl distilled water and 1M CH<sub>3</sub>CO<sub>2</sub>K.

### 2.3.4 Determination of Ferric Reducing Antioxidant Power (FRAP)

The reducing potential of each of the *Pupuru* flour blend extracts was determined by assessing the ability of the extract to reduce a FeCl<sub>3</sub> solution as described by Pulido et al. [20]. A 2.5 mL aliquot was mixed with 2.5 mL, 200 mM sodium phosphate buffer (pH 6.6) and 2.5 mL, 1% potassium ferricyanide. The mixture was incubated at 50°C for 20 min, and then 2.5 mL, 10% trichloroacetic was added and centrifuged at 650 g for 10 min. A 5 mL of the supernatant was mixed with an equal volume of water and 1 mL, 0.1% ferric chloride. The same treatment was performed to a standard ascorbic acid solution and the absorbance taken at 700 nm. The reducing power was then calculated and expressed as ascorbic acid equivalent.

$$\text{FRAP (mg/g)} = \frac{\text{Abs.sample} \times \text{Conc.standard}}{\text{Abs.standard} \times \text{Conc.sample}} \quad (4)$$

Where,

Abs<sub>standard</sub> = Absorbance of standard (Vitamin C),

Abs<sub>sample</sub> = Absorbance of reacting mixture with the test sample,

Conc<sub>standard</sub> = Stock concentration of standard in mg/ml, and

Conc<sub>sample</sub> = Stock concentration of sample in g/ml.

### 2.3.5 Determination of Total Phenol Content (TPC)

The total phenol content (TPC) was determined by Folin–Ciocalteu assay [21] using gallic acid as standard. Fifty microliters (50 µl) of aqueous extract solution containing 0.5 mg of aqueous extract was dispensed into a test tube, 50 µl of distilled water and 500 µl of Folin–Ciocalteu reagent was added respectively into the test tube and shaken thoroughly. After 3 min, 400 µl of 7.5% sodium carbonate solution was added and the mixture was incubated at 45°C in a water bath for 40 min. Absorbance was measured at 765 nm against the blank. The same procedure was repeated to all standard gallic acid solution (0.1 mg/ml). The blank is a mixture of 100 µl of distilled water, 500 µl of Folin–Ciocalteu reagent and 400 µl of 7.5% sodium carbonate. The total phenolic content was expressed as gallic acid equivalent per gram of sample (mg of GAE/g sample) through the calibration curve of gallic acid and calculated as follows;

$$\text{Total phenolic content (mg GAE/g)} = \frac{\text{Abs.sample} \times \text{Conc.standard (mg/ml)}}{\text{Abs.standard} \times \text{Conc.sample (mg/g)}} \quad (5)$$

## 2.4 Toxic and Phytochemical Content Determination

### 2.4.1 Determination of toxic property

#### 2.4.1.1 Determination of cyanide content

Some 50 mg of the sample was weighed out into a small flat-bottomed plastic vial [22]. Phosphate buffer (0.5 ml of 0.1 M at pH 4–10) was added, followed by the exogenous enzyme. A picrate paper attached to a plastic backing strip was added and the vial immediately closed with a screw stopper. After about 16 h at 30°C, the picrate paper was removed and immersed in 5.0 ml water for not less than 30 min. The absorbance was measured at 510 nm and the total cyanide content (ppm) determined by the equation:

$$\text{Total Cyanide content (ppm)} = \frac{396 \times \text{absorbance} \times 100}{z} \quad (6)$$

Where,

z = weight (mg) of sample.

## 2.4.2 Phytochemical

### 2.4.2.1 Determination of tannin content

Tannin content of the flour blends was determined using the method of Medoua et al. [23]. Two grams (2 g) of each sample was weighed into a 250 ml flask followed by addition of 200 ml of 0.004 M  $K_3Fe(CN)_6$  and 10 ml of 0.008 M  $FeCl_3$  in 0.008 M HCl. The flask was allowed to stand for 20 minutes and stirred occasionally at 10 min interval and 1 ml aliquot was removed. This aliquot was added 2 ml of 0.008 M  $FeCl_3$  in 0.008M HCl and 10 ml of 0.0015 M  $K_3Fe(CN)_6$ . After adding the final reagent, the absorbance was then read at 720 nm after 30 seconds against a blank.

$$Tannin (mg/g) = \frac{\text{Absolute of the sample} \times \text{concentration of the standard} \times \text{Dilution factor}}{\text{Absorbance of the standard} \times \text{sample size}} \quad (7)$$

### 2.4.2.2 Determination of phytate content

The determination of phytate in the sample was done using the method described by Abulude [24]. Eight grams (8 g) of each *Pupuru* flour blend was dispersed in 200 ml of 2% HCl and extracted. Following extraction, the dispersion was filtered and 50 ml of the filtrate was mixed with 10 ml of 0.3% ammonium cyanide ( $NH_4SCN$ ) and diluted with 107 ml of distilled water. The extract was titrated against 0.00195 g/ml of Ferric chloride solution until a brownish yellow colour persisted. Phytate content was estimated with the expression:

$$Phytate Phosphorous = (\text{Iron equivalent} \times 1.95 \text{ g of titre}) \times 3.65 \text{ g} \quad (8)$$

### 2.4.2.3 Determination of saponin content

The method described by Obadoni and Ochuko [25] was used to determine the saponin content of the *Pupuru* flour blends. Twenty grams (20 g) of each sample was put into a conical flask and 100 cm of 20% aqueous ethanol was added. This was heated over a hot water bath for 4 hours with continuous stirring at about 55°C. The mixture was filtered and the residue re-extracted with another 200 ml 20% aqueous ethanol. The combined filtrate was concentrated to 40 ml with the water bath at about 90°C. The concentrate was transferred into a 250 ml separator funnel and 20 ml of diethyl ether was added and shaken

vigorously. The aqueous layer was recovered while the ether layer was discarded. This purification step was repeated. 60 ml of n-butanol was added; the combined butanol extract was washed twice with 10 ml of 5% aqueous sodium chloride. The remaining solution was heated in a water bath; after evaporation, the samples were dried in the oven to constant weight; the saponin content was calculated in mg/g.

### 2.4.2.4 Alkaloid content determination

Alkaloids in the samples were determined by the method described by Harborne [26]. Five grams (5 g) of the sample was soaked in 200 mL of 20% acetic acid in ethanol for 4 hours. The mixture was filtered and the filtrate was concentrated on a water bath to about three-quarter of the original volume. Concentrated ammonia solution was added dropwise to the extract to precipitate the alkaloids. The solution was allowed to settle, the precipitate filtered and weighed while the alkaloid content was calculated as below:

$$\text{Alkaloid content} = \frac{\text{Weight of dry residue}}{\text{Weight of sample}} \times 100 \quad (9)$$

### 2.4.2.5 Determination of oxalate content

The method described by Ukpabi and Ejidoh [27] was used. Two grams (2 g) of each *Pupuru* sample was digested with 10 ml 6 M HCl for one hour and made up to 250 ml in a volumetric flask. The pH of the filtrate was adjusted with concentrated  $NH_4OH$  solution until the colour of the solution changed from a salmon pink colour to a faint yellow colour. Thereafter, the filtrate was treated with 10 ml of 5%  $CaCl_2$  solution to precipitate the insoluble oxalate. The suspension was centrifuged at 2500 rpm, after which the supernatant was decanted. The precipitate was dissolved in 10 ml of 20% (v/v)  $H_2SO_4$  and the solution was made up to 300 ml. An aliquot (125 ml) was heated until near boiling point and then titrated against 0.05 M standardized  $KMnO_4$  solution to a faint pink colour which persisted for about 30 seconds after which the burette reading was taken and used to estimate the oxalate content.

$$Oxalate (mg/g) = \frac{(\text{titre value} \times \text{volume of } KMnO_4 \times \text{dilution factor})/5}{\text{Sample size}} \quad (10)$$

## 2.5 Determination of Functional Properties

### 2.5.1 Foaming and emulsification capacities (%)

Foaming, emulsification, Water and oil-absorbing capacities (%) were determined using the methods of AOAC [28].

### 2.5.2 Swelling capacity

The swelling capacity of each sample was determined by the method described by [29]. One gram of flour was weighed into a 50 ml centrifuge tube. 50 ml of distilled water was added and mixed gently. The slurry was heated in a water bath at 90°C for 15 minutes. During heating, the slurry was stirred gently to prevent clumping of the flour. On completion, the tube containing the paste was centrifuged at 3,000 rpm for 10 minutes using a centrifuge machine. The supernatant was decanted immediately after centrifuging. The weight of the sediment was taken and recorded. The moisture content of sediment gel was thereafter determined to get the dry matter content of the gel.

$$\text{Swelling capacity} = \frac{\text{Weight of mass of sediment}}{\text{Weight of dry matter in the gel}} \quad (11)$$

### 2.5.3 Bulk density

Bulk density was determined using the method described by Wang and Kinsella [30]. 10 g of sample was weighed into a 50 ml graduated measuring cylinder. The sample was packed by gently tapping the cylinder on the benchtop. The volume of the sample was recorded.

$$\text{Bulk density} = \frac{\text{Weight of the sample}}{\text{The volume of the sample after tapping}} \quad (12)$$

### 2.5.4 Dispersibility (%)

Dispersibility of the samples was determined by the method described by Kulkarni et al. [31]. 10 g of flour was suspended in a 100 ml measuring cylinder and distilled water was added to reach a volume of 100 ml. The mixture was stirred vigorously and allowed to settle for 3 hours. The volume of settled particles was recorded and subtracted from 100. The difference was reported as percentage dispersibility of the sample:

$$\% \text{ dispersibility} = 100 - \frac{\text{the volume of the settled particle of sample}}{\text{total volume}} \quad (13)$$

### 2.5.5 Least gelation concentration

Least Gelation Concentration (LGC) of each sample was determined by the method described by Coffman and Garcia [32]. Ten suspensions (2, 4, 6, 8, 10, 12, 14, 16, 18 and 20% w/v) in 5 ml distilled water were prepared in test tubes. The test tubes containing the suspensions were heated in a boiling water bath (Thelco, model 83, USA) for 1 hour. The tubes and contents were cooled rapidly under running cold water and then cooled further for 2 hr at 4°C. The tubes were inverted to observe if the contents would fall or slip off. The least gelation concentration was that concentration when the sample from the inverted test tube did not fall or slip off.

## 2.6 Statistical Analysis

All the data obtained in this study were subjected to Analysis of Variance (ANOVA) using IBM SPSS version 21. Duncan Multiple Range Tests (DMRT) were carried out for the separation of means and determination of significant differences between means. Results are presented as mean  $\pm$  standard deviation accepted at  $P = .05$  confidence limit [33].

## 3. RESULTS AND DISCUSSION

### 3.1 In-vitro Antioxidant Activities

The in-vitro antioxidant activities of commercial and spontaneously-fermented African yam bean (AYB) (*Sphenostylis stenocarpa*) seed-enriched *Pupuru* flour blends are presented in Table 2. The observed trend was an increase in all the antioxidant activities as the levels of AYBS enrichment increased. The commercial and laboratory control samples showed no significant difference ( $P = .05$ ) in all the antioxidant capacities except DPPH. The ABTS activities of all the flour blends were significantly different ( $P = .05$ ) ranging from 7.61 (CP) to 12.27% (EP15). In the DPPH assay, radicals in the reaction medium were scavenged, resulting in a colour change from purple to yellow (indicating hydrogen donation ability of the sample), which led to a decrease in absorbance. The degree of solution discoloration, therefore, indicated the scavenging efficiency of the sample beverage [34]. The results of DPPH for all the samples were significantly different ( $P = .05$ ) between a range of 26.34% (CP) and 48.26% in EP15. The flavonoid in-vitro antioxidant capacities of all the flour blends ranged from 0.10 (CP) to 0.25 (mg CAE/g) (EP15). Flavonoids are a group of phenolic compounds extensively found in plants. As discovered with other biologically active anti-

nutrient compounds, flavonoids may stimulate beneficial and adverse physiological properties in humans [35]. Millet has been reported to contain flavonoids that possess a strong inhibitory consequence on thyroperoxidase activity in goitre cases which were caused by consumption of peanuts [35]. The healthy properties of flavonoids could be due to their antioxidative characteristics as free-radical neutralisers. Though, some particular functions such as cancer prevention, anti-inflammatory and antiviral activities, positive effect on capillary fragility and vascular protection have been documented [35]. Ferric reducing antioxidant power (FRAP) assay is used to determine the electron-donating abilities of natural antioxidants. FRAP of the entire samples ranged between 0.81 (P100) to 2.36 (mg/g) (EP15). Polyphenols are plant secondary metabolites that are abundant in plant-based foods. Total phenolic in-vitro antioxidant capacities of all the *Pupuru* flour blends varied from 15.74 (CP) to 24.15 mg GAE/g (EP15). Plant phenolic compounds are of utmost interest to researchers because of their antioxidant potential to prevent degenerative diseases [36]. The consumption of plants rich in phytochemicals has been reported to have a positive regulatory effect in humans and that phenolics possess the ability to scavenge free radicals, which normally accumulate in the body due to an imbalance between the antioxidant system of the body and the formation of reactive oxygen species [37]. The addition of AYBS to cassava in producing *Pupuru* flour blends in this study increased the phenolic contents of the samples to significant ( $P = .05$ ) levels, hence, their consumption could delay the ageing process, prevent degenerative diseases and enhance health.

### 3.2 Toxic Property and Phytochemical Characterisation

The toxic (cyanide) and phytochemical characterisation of commercial and spontaneously-fermented African yam bean (AYB) (*Sphenostylis stenocarpa*) seed-enriched *Pupuru* flour blends are as presented in Table 3. All the *Pupuru* flour blends contained significantly different ( $P = .05$ ) cyanide ranging from 0.87 - 2.51 mg/kg in EP15 and CP, respectively. The results indicated a reduction in cyanide content as the level of inclusion of AYBS increased. Cyanide intoxication can either be by inhalation or ingestion [38] and its level in cassava has been reported by Hahn [39] to depend on the cultivar, growth and environmental conditions,

such as temperature, soil type, humidity and age of the plant [40,41]. Cyanide has been reported as the major toxin that limits the consumption and utilisation of cassava roots which contains a range of 10 to 500 mg cyanide equivalents/kg DM [15]. Some cassava varieties, especially, the bitter one, contain more than 10mg cyanide equivalents/kg DM but the recommended limit to prevent acute toxicity in humans is  $< 10$  mg cyanide equivalents/kg DM [42]. The samples in this study are therefore, safe for consumption in terms their cyanide contents which are much lower than 10 mg/kg. Cyanide causes acute toxicity in humans, its residues (glucoside, cyanohydrin or free cyanide) in processed cassava, are as toxic as the unprocessed derivatives present in raw cassava roots but various effective detoxification methods which are usually in sequence and time-dependent have been documented to include peeling, grating, soaking, boiling/cooking, drying and fermentation [15,43-45].

The phytochemicals such as tannins, oxalates, phytates, are natural or synthetic compounds found in all plant foods (in varying quantities and types), which reduce absorption and utilisation of essential nutrients like vitamins and minerals. The results indicated that all the phytochemicals except tannins increased significantly ( $P = .05$ ) as levels of enrichment with AYBS increased.

Tannin contents were significantly different ( $P = .05$ ) ranging from 1.46 (in EP15) to 2.87 (mg/g) (in CP) in all the *Pupuru* flour blends. Tannins affect the nutritive values of food products, form a complex with protein (both substrate and enzyme) and inhibit digestion and absorption of proteins. Tannins also bind Iron, inhibit nonheme-Fe absorption and make it unavailable for utilization by the body [46]. A report [47] documented that condensed tannins may cleave to DNAs in the presence of copper ions, thereby, increasing malnutrition rate. Presence of tannins in large quantities in foods could lead to intestinal tract damage, and carcinogenesis [48]. Phytate in all the samples significantly ( $P = .05$ ) ranged from 0.85 (P100) to 1.40 mg/100 g (EP15), with the CP containing a lower value (1.12 mg/100 g) than in the laboratory control sample. The difference in these results might be due to different varieties used for both samples. Phytate possesses anti-nutritional activities in human diets due to its strong ability to chelate zinc, calcium and iron to form insoluble complexes which are not absorbed, hence, contributing to zinc and iron deficiencies [49]. Conversely,

phytate has a beneficial antioxidant property which subdues Fe-mediated ·OH formation by Fe, that is complexed by phytate [50] and acts as an anticancer agent [51]. A report [35] revealed that a focus has been on the beneficial properties of phytates owing to their antioxidative characteristics, which are favourable to offset free-radical activities. Saponins in all the samples were significantly ( $P = .05$ ) different, ranging between 4.18 (P100) and 13.27 mg/g (EP15), the commercial control (CP) sample had (5.97 mg/g) which was higher than what was obtained in the laboratory control (P100) sample. Saponins are compounds formed by triterpenoids or steroidal aglycones and a carbohydrate moiety by ester or ether linkages. They exist in diverse classes of plants, predominantly legumes, roots, and selected medicinal herbs. Their existence in food products has been considered to be harmful if consumed often, hence, their continued use in diets may compromise nutrient absorption [51]. Conversely, it has been claimed that they could also be beneficial since they display the ability to lower plasma cholesterol, they have anticancer activity, and they may act as an inhibitor of viral replication. It is not yet clear, though, whether the net effect in the diet would be negative [35]. Saponins have been reported to possess some bioactive components which are responsible for

metabolic and potential health benefits, treatment of various ailments, such as inflammation and fatigue [15]. They also provide energy, improve cognitive function and erectile dysfunction in men, and act on the central nervous system of humans to provide therapeutic effects.

The Alkaloid contents (mg/g) of the samples were significantly ( $P = .05$ ) different ranging from 24.89 (P100) to 29.05 mg/g (EP15), with the CP sample (27.78 mg/g) higher than the laboratory control sample. The variation in the alkaloid contents of both control samples might be due to differences in the varieties, genetic compositions and environmental factors of the cassava used. The observed trend in the alkaloid contents was an increase with increased enrichment levels of AYBS. Oxalates in the flour blends were also significantly ( $P = .05$ ) different ranging from 1.71 (P100) to 3.23 mg/g (EP15), with both the commercial and laboratory controls not significantly ( $P = .05$ ) different. The results for phytochemicals components indicated that enrichment of *Pupuru* flour blends with AYBS at 5%, 10% and 15% reduced cyanide and tannin which could be detrimental to consumers but increased the phytates, saponins, alkaloids and oxalates that might be of beneficial health implications.

**Table 2. In-vitro antioxidant activities of commercial and spontaneously-fermented African yam bean (AYB) (*Sphenostylis stenocarpa*) seed-enriched *Pupuru* flour blends**

Sample code	ABTS (%)	DPPH (%)	Flavonoid (mg CAE/g)	FRAP (mg/g)	Total Phenols (mg GAE/g)
CP	7.61±1.04 <sup>d</sup>	26.34±1.60 <sup>e</sup>	0.10±0.02 <sup>d</sup>	0.78±0.04 <sup>d</sup>	15.74±0.09 <sup>d</sup>
P100	7.97±1.32 <sup>d</sup>	29.41±1.37 <sup>d</sup>	0.10±0.03 <sup>d</sup>	0.81±0.02 <sup>d</sup>	16.30±0.13 <sup>d</sup>
EP5	9.19±0.84 <sup>c</sup>	37.95±2.43 <sup>c</sup>	0.15±0.02 <sup>c</sup>	1.84±0.03 <sup>c</sup>	17.14±0.80 <sup>c</sup>
EP10	10.94±1.02 <sup>b</sup>	43.21±1.69 <sup>b</sup>	0.19±0.01 <sup>b</sup>	2.27±0.06 <sup>b</sup>	21.47±1.01 <sup>b</sup>
EP15	12.27±0.96 <sup>a</sup>	48.26±1.48 <sup>a</sup>	0.25±0.02 <sup>a</sup>	2.36±0.01 <sup>a</sup>	24.15±0.62 <sup>a</sup>

Values are means ± SD with different superscripts in the same column are significantly different ( $P = .05$ ),  $n = 3$ . CP = Commercial 100% Cassava *Pupuru* Sample; P100 = Laboratory 100% Cassava *Pupuru*; EP5 = 5% AYB-enriched *Pupuru*; EP10 = 10% AYB-enriched *Pupuru*; EP15 = 15% AYB-enriched *Pupuru*

**Table 3. Toxic and phytochemical characterisation of commercial and spontaneously-fermented African yam bean (AYB) (*Sphenostylis stenocarpa*) seed-enriched *Pupuru* flour blends**

Sample code	Cyanide (mg/kg)	Tannin (mg/g)	Phytate (mg/ 100g)	Saponin (mg/g)	Alkaloids (mg/g)	Oxalate (mg/g)
CP	2.51±0.02 <sup>a</sup>	2.87±0.02 <sup>a</sup>	1.12±0.01 <sup>b</sup>	5.97±0.02 <sup>d</sup>	27.78±0.02 <sup>c</sup>	1.72±0.02 <sup>d</sup>
P100	2.17±0.06 <sup>b</sup>	2.09±0.02 <sup>b</sup>	0.85±0.03 <sup>c</sup>	4.18±0.30 <sup>e</sup>	24.89±0.04 <sup>e</sup>	1.71±0.10 <sup>d</sup>
EP5	1.71±0.03 <sup>c</sup>	1.71±0.05 <sup>c</sup>	1.12±0.18 <sup>b</sup>	10.00±0.28 <sup>c</sup>	25.37±0.02 <sup>d</sup>	1.84±0.08 <sup>c</sup>
EP10	1.10±0.01 <sup>d</sup>	1.69±0.02 <sup>d</sup>	1.15±0.50 <sup>b</sup>	12.36±0.30 <sup>b</sup>	28.93±0.05 <sup>b</sup>	2.76±0.04 <sup>b</sup>
EP15	0.87±0.01 <sup>e</sup>	1.46±0.04 <sup>e</sup>	1.40±0.04 <sup>a</sup>	13.27±0.24 <sup>a</sup>	29.05±0.03 <sup>a</sup>	3.23±0.05 <sup>a</sup>

Values are means ± SD with different superscripts in the same column are significantly different ( $P = .05$ ),  $n = 3$ . CP = Commercial 100% Cassava *Pupuru* Sample; P100 = Laboratory 100% Cassava *Pupuru*; EP5 = 5% AYB-enriched *Pupuru*; EP10 = 10% AYB-enriched *Pupuru*; EP15 = 15% AYB-enriched *Pupuru*



### 3.3 Functional Properties of *Pupuru* Flour Blends

The functional properties of spontaneously fermented *Pupuru* flour blends are as shown in Table 4. Foaming capacity (FC) is a property of protein in food samples which could be a benefit in their solubility, capacity to incorporate air for swelling, thus, giving the end product a honeycomb structure as desired in baked products and ice cream [52-54]. The ability of food materials to foam vary with the types of protein, solubility and additional factors [55]. Good foaming ability is a function of the flexible protein molecule that could reduce surface tension, conversely, the poor foaming ability is due to highly ordered globular protein which is relatively difficult to denature by heat [56]. The FC of spontaneously fermented *Pupuru* flour blends ranged significantly ( $P = .05$ ) from 3.28 in P100 to 4.92% for both EP5 and EP 15 while CP had 9.84% which was twice the highest value. These values were higher than those reported by [57,58] for African breadfruit and Kidney bean/wheat flour blends, respectively. High emulsion capacity of food products indicates better flavour retention, mouth feel and taste [59] and flours with good ECs will be useful for the preparation of comminuted meat products and their analogues [53]. Emulsification Capacity (EC) varied between 54.00 (P100) and 61.22% (EP15) with 47.06% in CP which was lower than the laboratory-prepared control (P100). These values were higher than those reported by [57, 58] for African breadfruit kernel and seed and Kidney bean/wheat flour blends, respectively. The disparity in EC of flour blends has been reported to be as a result of differences in the globular protein contents [60]. The ability to absorb water is a very vital property of flours and starches used in food preparations [61]. Water Absorption Capacity (WAC) of food suggests the quantity of water accessible for gelatinisation and is an indication of the amount of water retained in its protein matrix [31,54]. The WAC also shows the level of granular integrity and defines the weakness of associative forces between the starch granules, which permits more accessible molecular surfaces for binding with water molecules [62]. WAC indicates some product characteristics such as bulkiness, consistency, moistness, starch retrogradation and staling [63]. The (WAC) of the *Pupuru* flour blends ranged significantly ( $P = .05$ ) from 3.88 g/ml in EP10 and EP15 to 4.07 g/ml in P100 and EP5, while the CP had 1.92 g/ml. These values were higher than those reported by [57,58,64] in studies on

African breadfruit kernel and seed; Kidney bean/wheat; and smoked-dried *Pupuru* flour blends, respectively. The trend was a significant ( $P = .05$ ) reduction in WAC with the increased percentage of substitution with AYB seed flour, which might be as a result of low availability of polar amino acids which have been reported as the primary sites for water interaction of proteins [65,53]. The ability of food materials to absorb water is occasionally ascribed to its proteins content [30]. Increase in WAC in food systems permits food processors to manipulate the functional properties of doughs [66]. The values obtained in this study were desirable and could imply that the protein quality of the AYB seed flour used was good and able to bind a large quantity of water as reported by [67]. Oil absorption capacity (OAC) is a pointer to the rate at which protein adheres to fat in food formulations [56,54]. Fat serves as a flavour retainer and improves the mouthfeel of food products [68]. Flours with high oil absorption capacities are known as excellent meat extenders [53]. The OAC of the spontaneously fermented *Pupuru* flour blends differed significantly ( $P = .05$ ) ranging between 1.94 (EP15) and 2.14 g/ml (P100, EP5 and EP10) with 1.92 g/ml for CP. These values were lower than those reported by Ikegwu et al. [61] for starch from improved cassava cultivars and 3.80% reported by Yusuf et al. [69] for Jack bean starch.

Swelling Index (SI) of starchy food is the degree of the ability of starch to imbibe water and swell [61] The swelling indices of the spontaneously fermented *Pupuru* flour blends varied significantly ( $P = .05$ ) from 3.08 in EP15 and 3.79 g/ml in P100 with CP having 2.78 g/ml. There was no significant difference ( $P = .05$ ) among the swelling indices of CP, EP5 and EP10 but the P100 was significantly higher ( $P = .05$ ) and EP15, significantly lower ( $P = .05$ ). The values obtained in this study were lower than those (5.49 to 6.92 %.) reported by Ikegwu et al. [61] in a study that converted different cassava cultivars into starch. The lower results might be due to the inclusion of AYB seed in the flour blends, as well as various preliminary treatments such as fermentation, toasting and drying that have been applied to the flour blends in this study. The lower swelling indices also suggested a more highly ordered arrangement in their granules, as reported that swelling index of granules depicts the magnitude of associative forces within them [70], hence, the higher the swelling index, the lower the associative forces within a starchy food system.

**Table 4. Functional properties of spontaneously fermented *Pupuru* flour blends and the commercial control sample**

Parameters	CP	P100	EP5	EP10	EP15
Foaming capacity (%)	9.84±0.13 <sup>a</sup>	3.28±0.17 <sup>d</sup>	4.92±0.02 <sup>c</sup>	4.92±0.01 <sup>c</sup>	4.92±0.59 <sup>c</sup>
Emulsification (%)	47.06±0.70 <sup>e</sup>	54.00±0.99 <sup>d</sup>	56.86±0.31 <sup>c</sup>	56.86±0.89 <sup>c</sup>	61.22±1.11 <sup>ab</sup>
Water Absorption Capacity (g/ml)	1.92±0.03 <sup>e</sup>	4.07±1.89 <sup>a</sup>	4.07±0.64 <sup>ab</sup>	3.88±0.48 <sup>c</sup>	3.88±0.53 <sup>c</sup>
Oil Absorption Capacity (g/ml)	2.02±0.11 <sup>a</sup>	2.14±0.09 <sup>a</sup>	2.14±0.09 <sup>a</sup>	2.14±0.01 <sup>b</sup>	1.94±0.11 <sup>c</sup>
Swelling Index (g/ml)	2.78±0.02 <sup>b</sup>	3.29±0.01 <sup>a</sup>	2.89±0.53 <sup>b</sup>	2.88±0.01 <sup>b</sup>	2.58±0.03 <sup>c</sup>
Packed Bulk Density (g/ml)	0.76±0.14 <sup>a</sup>	0.77±0.96 <sup>a</sup>	0.77±0.73 <sup>a</sup>	0.72±0.31 <sup>b</sup>	0.72±0.78 <sup>b</sup>
Loose Bulk Density (g/ml)	0.52±0.05 <sup>bc</sup>	0.53±0.11 <sup>c</sup>	0.56±0.09 <sup>b</sup>	0.56±0.09 <sup>b</sup>	0.56±0.01 <sup>b</sup>
Dispersibility (%)	47.18±0.13 <sup>ab</sup>	47.14±0.03 <sup>a</sup>	42.86±0.05 <sup>c</sup>	42.85±0.53 <sup>c</sup>	45.71±0.01 <sup>b</sup>
Least Gelation (%)	0.80±0.21 <sup>a</sup>	1.00±0.00 <sup>d</sup>	1.00±0.00 <sup>d</sup>	1.20±0.00 <sup>c</sup>	1.40±0.00 <sup>b</sup>

Mean ± SD with different superscripts in the same column are significantly different ( $P = .05$ ),  $n = 3$ . Key: CP = Commercial *Pupuru* Sample; P100 = Laboratory 100% *Pupuru*; EP5 = 5% AYB-enriched *Pupuru*; EP10 = 10% AYB-enriched *Pupuru*; EP15 = 15% AYB-enriched *Pupuru*

A significant decrease ( $P = .05$ ) in swelling index of the P100 was observed from 3.79 to 3.08-3.39 for the AYB seed-enriched samples, which might have been as a result of the presence of lipids in the AYB seed which might have reduced the swelling capacity of the *Pupuru* flour particles, as reported by Cheftel et al. [71]. This trend was in agreement with the findings of [72,73] and the reduced swelling index with increased percentage inclusion of AYB seed might be attributed to the reduced starch component in the enriched samples which could have reduced the absorption of water. The reduced swelling capacity was attributed to a high-fat content which might have reduced the ability of a mixture of wheat and peanut flours to bind water [74]. Swelling capacity has been implicated for a greater volume and more feeling of satiety per unit weight of fermented cassava products to consumers while a swelling index of a minimum of 3.0 has been recommended as the preference of consumers [75]. Bulk Density (BD) of flour is a function of its particle size and is inversely proportional to bulk densities (Loose and bulk) [55]. The BD indicates heaviness, greater compactness and is imperative in determining the packaging requirements and material handling of flours [76]. The structure of starch polymers influences bulk density thus, a loose structure of starch polymers could result in low bulk density [77]. Packed bulk densities (g/ml) of *Pupuru* flour blends varied from 0.72 (EP10 and EP15) to 0.77 g/ml (P100 and EP5), while CP had 0.76 g/ml. Loose bulk densities of the *Pupuru* flour blends ranged between 0.53 (P100) and 0.56 g/ml (EP5, EP10 and EP15) while CP

had 0.52 g/ml which was not significantly different ( $P = .05$ ) from P100 but was significantly ( $P = .05$ ) lower than all the AYB seed-enriched samples (EP5, EP10 and EP15). These results indicated that the inclusion of AYB seed increased the loose bulk density of the *Pupuru* flour blends. There were no significant differences ( $P = .05$ ) in the Packed and loosed bulk densities of the control and the enriched samples *Pupuru* flour blends, as observed by [73] in a study on Soy-Melon Gari production. The result might have been due to the starch content of the flour blends which tends to make the mixture less bulky and lighter [66,73]. However, the packed bulk densities were consistently higher than the Loosed bulk densities, implying that more quantity of the enriched *Pupuru* flour blends can be packed better than the same specific volume of the control sample, as reported by Fagbemi [78]. Dispersibility of flour is its ability to reconstitute in water. The higher it is, the better the reconstitution of the flour in water [31] but the dispersibility value for flour samples during the storage is usually relatively high [62], hence, the ease to reconstitute and produce doughs with fine consistencies when mixed with water. The dispersibility of the *Pupuru* flour blends varied significantly ( $P = .05$ ) from 47.14 in P100 to 45.71% in EP15. There was no significant difference ( $P = .05$ ) between the CP and P100; EP5 and EP10 but EP15 were significantly ( $P = .05$ ) higher than all the others. These values were lower than those reported (69 – 86%) for orange flesh sweet potato-sorghum-soy flour blend [77]. Gelling ability of a food product is

predisposed by the nature of its inherent proteins, starch and gums, and their interaction during heat treatment [79]. Least gelation properties of the *Pupuru* flour blends ranged from 0.80 (CP) to 1.40% (EP15). There were significant differences ( $P = .05$ ) among all the samples, CP (0.80%) was significantly ( $P = .05$ ) lower than P100 (1.00), with EP5 and EP10 were 1.20% which were significantly ( $P = .05$ ) lower than EP15 (1.40%). The low least gelation values could be attributed to the probable development of intermolecular hydrogen bonds between amylose molecules and other proteins present in the cooled gel samples [80]. The rate of gelatinisation of starch is determined by the rate of starch granule swelling [81]. In a study that enriched *Gari* with Soy-Melon observed a higher gel strength in the control sample and reported that protein enriched-samples contained a relatively lower number of starch granules as compared to the unenriched samples [73].

#### 4. CONCLUSION

This study showed that the utilisation of African yam bean (*Sphenostylis stenocarpa*) seed (AYBS) to enrich *Pupuru* increased its In-vitro antioxidant capacity and phytochemical constituents, reduced the toxic cyanide content enhanced the functional properties. Thus, its suitability for consumption as a nutraceutical to scavenge free radicals, delay the ageing process, prevent or combat cardiovascular and degenerating diseases, and enhance overall health. The result is an indication that these combinations of *Pupuru* flour blends will lead significantly to the expansion of African yam bean (*Sphenostylis stenocarpa*) seed use, thus, lessening its potential extinction but generating income to local farmers.

#### COMPETING INTERESTS

Authors have declared that no competing interests exist.

#### REFERENCES

- Godfray CJ, Beddington JR, Crute IR, Haddad L, Lawrence D, Muir JF, et al. Food security: The challenge of feeding 9 billion people. *Science* 2010;327(5967): 812-818.
- World Bank. The World Bank annual report: Year in review (English). Washington, DC: World Bank; 2008.
- Food and Agriculture Organization of the United Nations. (FAO). The state of food insecurity in the world. Communication Division, Food and Agriculture Organization of the United Nations, Viale delle Terme di Caracalla, 00153 Rome, Italy. 2009;1-61.
- Omar NF, Hassan SA, Yusoff UK, Abdullah NAP, Wahab PEM, Sinniah U. Phenolics, flavonoids, antioxidant activity and cyanogenic glycosides of organic and mineral-base fertilized cassava tubers. *Molecules*. 2012;17:2378-2387. DOI: 10.3390/molecules17032378
- Olaiya CO, Idowu PA, Karigidi KO. Antioxidative and antimicrobial activities of corn steep liquor anti-diabetic herb extracts. *Annals Food Science and Technology*. 2016;17(2); 272-279.
- Soetan KO, Olaiya CO and Karigidi KO. Comparative *In vitro* antioxidant activities of six accessions of African yam beans (*Sphenostylis stenocarpa* L.). *Annals. Food Science and Technology*. 2018;19(3):455-461.
- Yi B, Hu L, Mei W, Zhou K, Wang H, Luo Y et al. Antioxidant phenolic compounds of cassava (*Manihot esculenta*) from Hainan. *Molecules*. 2011;16(12):10157–10167. DOI: 10.3390/molecules161210157 [PMCID: PMC6264345] [PMID: 22157579]
- Giacomo CD, Acquaviva R, Sorrenti V, Vanella A, Grasso S, Barcellona ML, et al. Oxidative and antioxidant status in plasma of runners: Effect of oral supplementation with natural antioxidants. *J. Med. Food*. 2009;12:145–150. DOI: 10.1089/jmf.2008.0074
- Zhang Z, Liao L, Moore J, Wu T, Wang Z. Antioxidant phenolic compounds from walnut kernels (*Juglans regia* L.) food Chemistry. 2009;113:160–165. DOI: 10.1016/j.foodchem.2008.07.061
- Xu BJ, Chang SKC. A comparative study on phenolic profiles and antioxidant activities of legumes as affected by extraction solvents. *Journal of Food Science*. 2007;72(2):159-166.
- Granato D, Katayama FCU, de Castro IA. Phenolic composition of South American red wines classified according to their antioxidant activity, retail price and sensory quality. *Food Chemistry*. 2011;129:366–373. DOI: 10.1016/j.foodchem.2011.04.085
- Enujiugha VN, Talabi JY, Malomo SA and Olagunju AI. DPPH radical scavenging capacity of phenolic extracts from African

- yam bean (*Sphenostylis stenocarpa*). Food and Nutrition Sciences. 2012;3(1):7-13. DOI: 10.4236/fns.2012.31002
13. Charles AL, Ssiroth K, Huang TC. Proximate composition, mineral contents, hydrogen cyanide and phytic acid of 5 cassava genotypes, Food Chemistry. 2005;92(4):615-620.
  14. Ayaşan T. Use of cassava and products in animal nutrition (in Turkish). J Agric Fac Gaziosmanpasa University. 2010;27(1): 73-83.
  15. Montagnac JA, Davis CR, Tanumihardjo SA. Processing techniques to reduce toxicity and antinutrients of cassava for use as a staple food. Comprehensive Review in Food Science and Food Safety. 2009;8(1):17-27. Available:<https://doi.org/10.1111/j.1541-4337.2008.00064.x>
  16. Oluwamukomi MO, Akinlabi AA. Nutritional Enrichment of “Pupuru” and smoked-dried Cassava product with African yam bean Seed flour. Annals of Science and Biotechnology. 2011;2(1):26-35.
  17. Re R, Pellegrini N, Proteggente A, Pannala A, Yang M and Rice-Evans C. Antioxidant activity applying an improved ABTS radical cation decolorization assay. Free Radical Biology and Medicine. 1999;26:1231-37.
  18. Gyamfi MA, Yonamine M, Aniya Y. Free-radical scavenging action of medicinal herbs from Ghana: *Thonningia sanguinea* on experimentally-induced liver injuries. General Pharmacology Vascular System. 1999;32:661-667.
  19. Bushra IF, Abdul W, Ali R, Ullah H, Iqbal H, Almas, M, et al. Antimicrobial activity of *Malva neglecta* and *Nasturtium microphyllum*. International Journal of Research on Ayurveda Pharmacology. 1999;3:808-810.
  20. Pulido R, Bravo L, Saura-Calixto F. Antioxidant activity of dietary polyphenols as determined by a modified ferric reducing/antioxidant power assay. Journal of Agriculture and Food Chemistry. 2000; 48:3396-3402.
  21. Singleton VL, Orthofer R, Lamuela-Raventos RM. Analysis of total phenols and other oxidation substrates and antioxidants by means of Folin-Ciocalteu reagent. Methods Enzymology. 1999;299: 152-179.
  22. Egan SV, Yeoh HH, Bradbury JH. Simple picrate paper kit for determination of the cyanogenic potential of cassava flour. Journal of Science, Food and Agriculture. 1998;76:39-48.
  23. Medoua NG, Mbome Lape I, Agbor-Egbe T, Mbofung CMF. Antinutritional factors changes occurring in trifoliolate yam (*Dioscorea dumetorum*) tubers after harvest. Food Chemistry. 2007;102:716–20.
  24. Abulude FO. Effect of processing on nutritional composition, phytate and functional properties of Rice (*Oryza sativa* L) flour. Nigerian Food Journal. 2004;22: 97-104
  25. Obadoni BO, Ochuko PO. Phytochemical studies and comparative efficacy of the crude extract of some homeostatic plants in Edo and Delta states of Nigeria. Global Journal of Pure Applied Science. 2001;8: 203-208.
  26. Harborne JB. Phytochemical Methods, A Guide to Modern Techniques of Plant Analysis. Chapman and Hall, London. 1973; 113.
  27. Ukpabi A, Ejidoh EO. Experimental procedures for food and water analysis. San Press Publishers, Enugu, Nigeria. 1989;89.
  28. AOAC. Association of Official Analytical Chemist. Official Methods of Analysis of the Analytical Chemist International, 18<sup>th</sup> ed. Gathersburg, MD USA; 2012.
  29. Takashi S, Sieb PA. Paste and gel properties of prime corn and wheat starches with and without native lipids. Cereal Chemistry. 1988;65:474-83.
  30. Wang JC, Kinsella JE. Functional properties of novel proteins: Alfalfa leaf protein. Journal of Food Science. 1976;41: 286-92.
  31. Kulkarni KD, Kulkarni DN, Ingle UM. Sorghum Malted and soya bean weaning food formulations: Preparation, functional properties and nutritive value. Food and Nutrition Bulletin. 1991;13:322-27.
  32. Coffman CW, Garcia VV. Functional properties and Amino acid content of protein isolate from mug bean flour. Journal of Food Technology. 1977;12:473 – 84.
  33. Steel R, Torrie J and Dickey D. Principles and procedures of statistics: A biometrical approach, 3<sup>rd</sup> ed., McGraw Hill Book Co., New York, USA; 1997.
  34. Sánchez-Moreno C. Methods used to evaluate the free radical scavenging activity in foods and biological systems.

- Food Science and Technology International. 2002;8:121-37.
35. Trugo LC, von Baer E, von Baer D. Reference Module in Food Science. 2016. Elsevier Ltd. Available:<http://dx.doi.org/10.1016/B978-0-08-100596-5.00211-0>.
  36. Roginsky V. Chain-breaking antioxidant activity of natural polyphenols as determined during the chain oxidation of methyl linoleate in Triton X-100 micelles. *Archive of Biochemistry and Biophysics*. 2003;414:261-70.
  37. Savikin K, Zdunić G, Janković T, Tasić S, Menković N, Stević T, et al. Phenolic content and radical scavenging capacity of berries and related jams from certificated area in Serbia. *Plant Foods and Human Nutrition*. 2009;64:212-17.
  38. Jack B, France B, Venkatesh M and Peter R. Food fortification - the debate continues. *Science in Africa*. 2002;75. Available:<http://scienceinafrica.com/old/2002/june/food.htm>
  39. Hahn S. An overview of traditional processing and utilization of cassava in Africa. Cassava as Livestock Feed in Africa: Proceedings of the IITA/ILCA/University of Ibadan Workshop on the Potential Utilization of Cassava as Livestock Feed in Africa: 14-18, Ibadan, Nigeria; 1992.
  40. Food and Agriculture Organization (FAO) of the United Nations Rome. The state of food and agriculture. *FAO Agriculture Series*, no. 23; 1990. [ISBN: 92-5-102989-X]
  41. Paula AC, Estevao M, Mario E, Fernando M, Julie CM, Rezaul H. et al. Processing of cassava roots to remove cyanogen". *Journal of Food Composition and Analysis*. 2005; 18:451-60.
  42. Food and Agriculture Organization (FAO)/ World Health Organization (WHO). Joint FAO/WHO Food standard programme. *Codex Alimentarius XII Suppl*. 4 Ed. Rome. 1991;1-42.
  43. Lambri M, Fumi MD, Roda A, Marco DD. Improved processing methods to reduce the total cyanide content of cassava roots from Burundi. *African Journal of Biotechnology*. 2013;12(19):2685-91.
  44. Kyawt YY, Imai Y, Yara T, Kawamoto Y. Effect of ensiling process and additive effects of fermented juice of epiphytic lactic acid bacteria on the cyanide content of two varieties of cassava. *Animal Nutrition and Feed Technology*. 2014;14(3):447.
  45. Guira Flibert TA, Savadogo A. African cassava traditional fermented food: The microorganism's contribution to their nutritional and safety values - A review. *International Journal of Current Microbiology and Applied Sciences*. 2016;5 (10):664-87. DOI:<http://dx.doi.org/10.20546/ijcmas.2016.510.074>
  46. Adeniji TA, Sanni LO, Barimalaa IS, Hart AD. Anti-nutrients and heavy metals in some new plantain and banana cultivars. *Nigerian Food Journal*. 2007;25(1):171-77.
  47. Aletor V. Allelochemicals in plant foods and feeding stuffs: 1. Nutritional, biochemical and physiopathological aspects in animal production. *Veterinary Human Toxicology*. 1993;35:57-67.
  48. Anuonye JC, Jigam AA and Ndaceko GM. Effects of extrusion-cooking on the nutrient and anti-nutrient composition of pigeon pea and unripe plantain blends. *Journal of Applied Pharmaceutical Sciences*. 2012;2 (5)158-62.
  49. Raboy V. Seeds for a better future: 'low phytate' grains help to overcome malnutrition and reduce pollution. *Trends in Plant Science*. 2001;6(10):458-62.
  50. Graf E, Empson KL, Eaton JW. Phytic acid. A natural antioxidant. *J Biol Chem*. 1987;262(24):11647-50.
  51. Graf E, Eaton JW. Suppression of colonic cancer by diet-ary phitic acid. *Nutrition and Cancer*. 1993;19:11-19.
  52. Fenema OR. *Principle of food science. Food Chemistry*. Marcel Dekker Inc. New York. 2000;125:429.
  53. Adeyanju BE, Enujiugha VN and Bolade MK. Effects of addition of kidney bean (*Phaseolus vulgaris*) and alligator pepper (*Aframomum melegueta*) on some properties of 'aadun' (a popular local maize snack). *Journal of Sustainable Technology*. 2016;7(1):45-58.
  54. Akinola SA, Enujiugha VN. Physicochemical and sensory qualities of "aadun" a maize based snack supplemented with defatted African oil bean seed flour. *Applied Tropical Agriculture*. 2017;22(2):188-96.
  55. Akubor PI, Eze JI. Quality evaluation and cake making potential of sun and oven dried carrot fruit. *International Journal of Bioscience*. 2012;2(10):19 -27.

56. Onimawo IA, Akubor PI. Food chemistry (Integrated approach with biochemical background). 2<sup>nd</sup> edn. Joytal printing press, Agbowo, Ibadan, Nigeria; 2012.
57. Akubor PI, Isolokwu PC, Ugbane O, Onimawo A. Proximate composition and functional properties of African breadfruit kernel and flour blends. Food Research International. 2000;33(8):707-12.
58. Okoye JI, Nkwocha AC, Agbo AO. Chemical composition and functional properties of kidney bean/wheat flour blends. Journal of Food Science and Technology. 2008;2:27-32.
59. Oyarekua MA, Adeyeye EI. Comparative evaluation of the nutritional quality, functional properties and amino acid profile of co-fermented maize/cowpea and sorghum/cowpea Ogi as infant complementary food. Asian Journal of Chemistry and Nutrition. 2009;1:31-39
60. Sathe SK, Deshpande SS, Salunkhe DK. Functional properties of winged bean (*Psophocarpus tetragonolobus*, L) proteins. Journal of Food Science. 1982; 47:503-6.
61. Ikegwu OJ, Nwobasi VN, Odoh MO, Oledinma NU. Evaluation of the pasting and some functional properties of starch isolated from some improved cassava varieties in Nigeria. African Journal of Biotechnology. 2009;8(10):2310-15.
62. Adebowale YA, Adeyemi IA, Oshodi AA. Functional and physicochemical properties of flours of six *Mucuna* species. African Journal of Biotechnology. 2005;4(12): 1461-68.
63. Siddiq M, Ravi R, Harte JB, Dolan KD. Physical and functional characteristics of selected dry bean (*Phaseolus vulgaris* L.) flours. LWT – Food Science and Technology. 2010;43(2):232–37.
64. Shittu TA, Adedokun II. Comparative evaluation of the functional and sensory characteristics of three traditional fermented cassava products. Journal of Natural Sciences, Engineering and Technology. 2010;9(2):106-16.
65. Kuntz ID. Hydration of macromolecules III. Hydration of polypeptides Journal of American Chemical Society. 1971;3:514–15.
66. Iwe MO, Onadipe OO. Effect of addition of extruded full fat soy flour into sweet potato flour on functional properties of the mixture. Journal of Sustainable. Agriculture and Environment. 2001;3(1):109-17.
67. Giami SY, Bekebani DA, Emelike NJ. Proximate composition and functional properties of winged bean (*Psophocarpus tetragariolobus*) Nutrition Science. 1992;3: 36-38.
68. Balogun IO, Olatidoye OP. Functional properties of dehulled and undehulled velvet beans flour (*Macuna utilis*). Journal of Biological Science and Bioconservation. 2010;2:1-10.
69. Yusuf AA, Ayedun H, Logunleko GB. Functional properties of unmodified and modified Jack bean (*Canavalia ensiformis*) starches. Nigerian Food Journal. 2007;25 (2):50852.
70. Sanni LO, Maziya-Dixon B, Akanya CI, Alaya Y, Egwuonwu CV, Okechukwu RU, et al. Standards for cassava products and guidelines for export. International Institute of Tropical Agriculture, Ibadan, Nigeria. 2005;11-39.
71. Cheftel GS, Cuq JL, Loriet D. Amino acids, peptides and proteins. In: Food Chemistry, 2<sup>nd</sup> ed., Fennema, O.R, ed. Marcel Dekker, New York. 1985;245-369.
72. Oluwamukomi MO, Adeyemi IA, Oluwalana IB. Effects of soybean enrichment on the physicochemical and sensory properties of gari. Applied Tropical Agriculture. 2005;10(Special issue):44-49.
73. Oluwamukomi MO, Jolayemi OS. Physico-thermal and pasting properties of soy-melon-enriched “gari” semolina from cassava. Agricultural Engineering International: CIGR Journal. 2012;14 (3):105-16.
74. Prinyawiwatkul W, McWatters KH, Beuchart LR, Phillips RD. Physical properties of cowpea paste and akara as affected by supplementation with peanut flour. Journal of Agriculture and Food Chemistry. 1994;42:1750-56.
75. Almazan AM. Influence of cassava variety and storage on gari quality. Tropical. Agriculture (Trinidad). 1992;69(4):386-90.
76. Karuna D, Noel G, Dilip K. Food and Nutrition Bulletin. United Nation University. 1996;17:2.
77. Alawode EK, Idowu MA, Adeola AA, Oke EK, Omoniyi SA. Some quality attributes of complementary food produced from flour blends of orange flesh sweet potato, sorghum, and soybean. Croatia Journal of Food Science and Technology. 2017;9(2): 122–29.
78. Fagbemi TN. Effect of blanching and ripening on functional properties of

- Plantain (*Musa aab*) Flour. Plant Food for Human Nutrition. 1999; 54(2):261–69.
79. Enujiugha VN, Ayodele-Oni O. Evaluation of nutrients and some anti-nutrients in lesser-known, underutilized oilseeds. International Journal of Food Science and Technology. 2003;38:525–28.
80. Mbaeyi IE. Production and evaluation of breakfast cereal using pigeon pea (*Cajanus cajan*) and sorghum (*Sorghum bicolor* L.). M.Sc. in Food Science and Technology, Faculty of Agriculture, University of Nigeria, Nsukka. 2005; 167.
81. Zobel HF. Starch gelatinization and mechanical properties of starch pastes. Page 285 in: Starch: Chemistry and technology. RL Whistler, JN Bemiller, EF Paschall, eds. Academic Press: Orlando, FL; 1984.

---

© 2020 Isaac-Bamgboye et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

*Peer-review history:*

*The peer review history for this paper can be accessed here:*  
<http://www.sdiarticle4.com/review-history/55044>